arXiv:1608.06573v2 [math.CA] 2 Dec 2016

Standard transmutation operators for the one dimensional Schrödinger operator with a locally integrable potential

Hugo M. Campos
School of Mathematics, Yachay Tech, Yachay City of Knowledge,
100119, Urcuqui, Ecuador
email: hfernandes@yachaytech.edu.ec

December 5, 2016

Abstract

We study a special class of operators T satisfying the transmutation relation

$$(\frac{d^2}{dx^2} - q)Tu = T\frac{d^2}{dx^2}u$$

in the sense of distributions, where q is a locally integrable function, and u belongs to a suitable space of distributions depending on the smoothness properties of q.

A method which allows one to construct a fundamental set of transmutation operators of this class in terms of a single particular transmutation operator is presented. Moreover, following [27], we show that a particular transmutation operator can be represented as a Volterra integral operator of the second kind.

We study the boundedness and invertibility properties of the transmutation operators, and use these to obtain a representation for the general distributional solution of the equation $\frac{d^2u}{dx^2} - qu = \lambda u$, $\lambda \in \mathbb{C}$, in terms of the general solution of the same equation with $\lambda = 0$.

Keywords: Transmutation, Transformation operator, Schrödinger operator, Goursat problem, Weak solution, Spectral parameter power series.

1 Introduction

If L and M are differential operators, then an operator T satisfying the relation $LT = TM$, in a suitable functional space, is called a transmutation operator. Transmutation operators are a useful tool in the study of differential operators with variable coefficients, see [1], [9], [25], [26], and [27] for their classical theory.

In the present paper we deal mainly with transmutations corresponding to the pair $L = \frac{d^2}{dx^2} - q(x)$ and $M = \frac{d^2}{dx^2}$. It is well known that if $q \in C[-a,a]$, $a > 0$, then a transmutation operator for such pair on $C^2[-a,a]$ can be represented in the form

$$\mathbf{T}u(x) = u(x) + \int_{-x}^{x} K(x,t)u(t)dt, \tag{1}$$

where $K(x,t)$ is the unique solution of a certain Goursat problem [27].

Recent work [8] shows an interesting connection between the transmutation operators, and the concept of L-base introduced in [11]. There, a parametrized family of operators $\{T_c\}$ was introduced possessing the following properties: *i)* for each $c \in \mathbb{C}$, T_c is a Volterra integral operator and $T_0 = \mathbf{T}$; *ii)* if $q \in C^1[-a,a]$, then T_c is a transmutation operator on $C^2[-a,a]$ for suitable values of c; *iii)* the family of functions $\{\varphi_k\}$, where $\varphi_k = T_c[x^k]$, $k = 0,1,2,..$, is an L-base.

It is worth mentioning that the L-base $\{\varphi_k\}$ is the main ingredient to obtain the so called spectral parameter power series (SPPS) representation, a useful representation of the general solution of the Sturm–Liouville equation [17], [19], and it also arises in Bers's theory of pseudoanalytic functions [3], [18] in the study of a special kind of Vekua equations. These connections have led to new applications of the transmutation operators to the study of different problems arising in mathematical physics [7], [8], [13], [14], [20], [24].

Regarding these applications, it is desirable to have the transmutation property of T_c under the most general scenario, which, to the best of our knowledge, is valid for a continuous potential q and for all parameters $c \in \mathbb{C}$ (see Remark 24 for further details).

The first goal of this paper is the study of transmutations for the Schrödinger operator with integrable coefficients. Our approach is not based on merely extending existing results from continuous coefficients, but instead we make our study in the spirit of standard L-bases (see Definition 7). We will consider the class of transmutation operators, here named standard, which have the property of mapping the nonnegative integer powers of the independent variable into an s-L-base. This class includes the transmutation operators having the form of a Volterra integral of the second kind, and others.

This paper is organized as follows. In Section 2 we fix some notation and we give some auxiliary results needed throughout the paper.

In Section 3, based on the properties of the s-L-bases, we arrive at several results concerning the s-transmutation operators, and we establish the first main result of the paper, Theorem 10, that if a bounded operator T on $L^1[a,b]$ maps the powers of the independent variable into an s-L-base, then T is a transmutation operator on $W^{2,1}[a,b]$. This result represents another proof for the transmutation property of the operator T_c and also suggests that some properties or relations on the s-transmutations might be studied first on smaller spaces, like the linear space of polynomials $\mathcal{P}(\mathbb{R})$, and then extended to suitable Banach spaces.

In Section 4 we present a fairly simple method for constructing new transmutation operators on $\mathcal{P}(\mathbb{R})$ when one single s-transmutation is known (Theorem 17) and we also provide explicit formulas relating any two s-transmutations on $\mathcal{P}(\mathbb{R})$, allowing the construction of one of them in terms of the other (Proposition 18).

From the viewpoint of applications, it is desirable to have the transmutation property (5) on larger spaces than $\mathcal{P}(\mathbb{R})$. Section 5 is devoted to the study of s-transmutations on Sobolev spaces. Extending the results of [27], we show that a particular s-transmutation operator on $W^{2,1}[-a,a]$ can be represented in the form (1), where $K(x,t)$ is a weak solution of a Goursat problem (Theorem 21). Combining this result with those of Section 4, we arrive at the main result of the paper, Theorem 22, an explicit representation of the general s-transmutation operator on Sobolev space. As an application, we get the SPPS representation for the general solution of the equation $\frac{d^2u}{dx^2} - qu = \lambda u$ in a slightly more general setting than was known previously (Corollary 23).

Even in the case of continuous coefficients, the transmutation relation (5) has been studied mainly in the space of twice differentiable functions, and assuming the existence of all the involved derivatives in the classical sense. However, as always, for the best understanding of differential operators, the framework of distributions is the most suitable.

The second goal of the paper is to study the s-transmutations on spaces of distributions, which is done in Section 6. Assuming that the potential of the Schrödinger operator is locally integrable, we show that each s-transmutation operator admits a continuous extension as a linear operator and as a transmutation operator to a suitable space of distributions, depending on the smoothness of the potential q. Also, this procedure leads to the construction of all the distributional s-transmutation operators.

Lastly, in Section 7 we provide some conclusions and generalizations.

2 Preliminaries

First, following [2], [5], and [28], let us introduce the spaces we are going work with. Let $(a,b) \subset \mathbb{R}$ be a bounded open interval and denote by $F(a,b)$ any one of the following classical spaces: $L^p(a,b)$, $C^k(a,b)$, or $C^\infty(a,b)$, for $k \in \mathbb{N}_0 = \{0,1,2..\}$ and $1 \leq p \leq \infty$. Note that $L^p(a,b)$ and $C^k(K)$ are Banach spaces with the corresponding natural norms whenever $K \subset (a,b)$ is compact, and $C^\infty(K)$ is a Fréchet space when equipped with the topology of uniform convergence on K in each derivative [2].

Let us denote by $F_{loc}(a,b)$ the subspace of $F(a,b)$ formed by those functions $\psi \in F(K)$, for each compact set $K \subset (a,b)$. By $F_c(a,b)$ we denote the set of functions $\psi \in F(a,b)$ such that $\operatorname{supp}\psi \subset (a,b)$, here $\operatorname{supp}\psi$ is defined as the complement of the largest open set on which ψ vanishes a.e. We endow $F_c(a,b)$ with the inductive limit topology, turning $F_c(a,b)$ into a locally convex topological vector space [2], [28]. A sequence of functions $\psi_n \in F_c(a,b)$ converges to $\psi \in F_c(a,b)$ with respect to this topology if there exists a compact set $K \subset (a,b)$ such that $\operatorname{supp}\psi_n \subset K$ for all n and $\psi_n \to \psi$ in $F(K)$ as $n \to +\infty$. Let $(F_c(a,b))'$ be the dual space of $F_c(a,b)$, consisting of all continuous functionals $u : F_c(a,b) \to \mathbb{C}$. The value of a functional u on $\psi \in F_c(a,b)$ is denoted by $\langle u, \psi\rangle$ and continuity means that $\langle u, \psi_n\rangle \to \langle u, \psi\rangle$ provided that $\psi_n \to \psi$ in $F_c(a,b)$ as $n \to \infty$. Convergence in $(F_c(a,b))'$ is defined to be pointwise convergence. Obviously, for a smaller set of $F_c(a,b)$, the set $(F_c(a,b))'$ will be bigger.

The space $C_c^\infty(a,b)$, called the space of test functions, plays an important role in the

theory of distributions. As defined above, $\phi_n \to \phi$ in $C_c^\infty(a,b)$ if there exists a compact set $K \subset (a,b)$ such that $\operatorname{supp}\phi_n \subset K$ for all n and $\frac{d^m\phi_n}{dx^m} \to \frac{d^m\phi}{dx^m}$ uniformly on K as $n \to \infty$ for each $m \in \mathbb{N}_0$. A distribution is an element from the dual of $C_c^\infty(a,b)$, which from now on we denote by $\mathcal{D}'(a,b)$, i.e., $\mathcal{D}'(a,b) \equiv (C_c^\infty(a,b))'$. A distribution u is called regular if there exists an $f \in L^1_{loc}(a,b)$ such that $\langle u,\phi\rangle = \int_a^b f(x)\phi(x)dx$ for all $\phi \in C_c^\infty(a,b)$. We usually identify a regular distribution u with the corresponding function f, $u \equiv f$.

Definition 1 *Let T and $T^\square$ be linear operators from $C_c^\infty(a,b)$ to $L^1_{loc}(a,b)$ such that $\int_a^b (T\psi)\phi dx = \int_a^b \psi(T^\square\phi)dx$ for all $\psi,\phi \in C_c^\infty(a,b)$. The operator $T^\square$, if it exists, is called the transpose of T.*

The most common way to extend linear operations from functions to distributions is the following.

Proposition 2 *Let T be a linear operator defined on $C_c^\infty(a,b)$. If $T^\square : C_c^\infty(a,b) \to C_c^\infty(a,b)$ is continuous, then the operator T can be extended to a continuous linear operator from $\mathcal{D}'(a,b)$ to itself by the formula*

$$\langle Tu,\phi\rangle := \left\langle u, T^\square\phi\right\rangle \quad , u \in \mathcal{D}'(a,b), \phi \in C_c^\infty(a,b).$$

Proof. This is a special case of [12, Th. 3.8]. ■

As $\frac{d}{dx}^\square = -\frac{d}{dx}$ is continuous on $C_c^\infty(a,b)$, the derivative of $u \in \mathcal{D}'(a,b)$ is defined by employing the previous proposition: $\langle u',\phi\rangle := -\langle u,\phi'\rangle$. It is again an element of $\mathcal{D}'(a,b)$.

The Sobolev space $W^{k,p}(a,b)$, $k \in \mathbb{N}$, $1 \le p \le \infty$ consists of all functions $f \in L^p(a,b)$ having distributional derivatives $f^{(n)} \in L^p(a,b)$ up to order $n \le k$.

Denote the set of complex-valued absolutely continuous functions on $[a,b]$ by $AC[a,b]$. We recall that a function V belongs to $AC[a,b]$ iff there exists $v \in L^1(a,b)$ and $x_0 \in [a,b]$ such that $V(x) = \int_{x_0}^x v(t)dt + V(x_0), \forall\ x \in [a,b]$ (see, e.g., [2], [16]). Thus, if $V \in AC[a,b]$, then $V \in C[a,b]$ and V is a.e. differentiable with $V'(x) = v(x)$ on (a,b). Moreover, the usual derivative of a function from $AC[a,b]$ coincides a.e. with its distributional derivative [16], thus $AC[a,b] \subseteq W^{1,1}(a,b)$. In fact, the linear spaces $W^{1,1}(a,b)$ and $AC[a,b]$ coincide (see [5], Theorem 8.2) in the following sense: $u \in W^{1,1}(a,b)$ iff there exists $V \in AC[a,b]$ such that $u = V$ a.e. on (a,b).

Consider the differential equation

$$u'' + qu = h, \tag{2}$$

in the sense of distributions, where $q,h \in L^1(a,b)$ and the distribution qu is defined by $\langle qu,\varphi\rangle := \langle u, q\varphi\rangle$. Note that if $q \notin C^\infty(a,b)$, then it might happen that $u \in \mathcal{D}'(a,b)$ but $qu \notin \mathcal{D}'(a,b)$. To overcome this situation, we first will look for solutions of (2) in $W^{1,1}(a,b)$, which are called weak solutions.

Proposition 3 *Let q and h be functions in $L^1(a,b)$. Then a function $u \in W^{1,1}(a,b)$ is a weak solution of (2) iff $u \in W^{2,1}(a,b)$ and (2) is satisfied a.e. on (a,b).*

Proof. See [4]. ■

The last result reveals the density of polynomials in $W^{2,1}(a,b)$.

Proposition 4 *Let $u \in W^{2,1}(a,b)$. Then there exists a sequence of polynomials P_n such that $P_n \to u$, $P_n' \to u'$ uniformly on $[a,b]$ and $P_n'' \to u''$ in $L^1(a,b)$, as $n \to \infty$.*

Proof. Let $u \in W^{2,1}(a,b)$. Then $u'' \in L^1(a,b)$, and because polynomials are dense in $L^1(a,b)$ there exists a sequence of polynomials Q_n such that

$$Q_n \to u'' \text{ in } L^1(a,b), \text{ as } n \to \infty. \tag{3}$$

Integrating (3) twice we see that the sequence of polynomials $P_n(x) := u(0) + u'(0)x + \int_0^x \int_0^t Q_n(s)\, ds\, dt$ fulfills the conclusion of the proposition. ■

3 s-transmutations and L-bases. First results

Consider the one-dimensional Schrödinger operator

$$L = \frac{d^2}{dx^2} - q(x), \tag{4}$$

with domain of definition $D_L = W^{2,1}(a,b)$, where $q \in L^1(a,b)$.

Definition 5 *A linear map T is called a transmutation operator on $\chi \subseteq D_L$ if for every $u \in \chi$ we have $u'' \in D_T$, $Tu \in D_L$ and the following equality holds*

$$\left(\frac{d^2}{dx^2} - q\right) Tu = T\frac{d^2}{dx^2}u. \tag{5}$$

Let $\mathcal{P}(\mathbb{R})$ denote the linear space of polynomials. We will be mainly interested in operators satisfying the transmutation property (5) on sets $\chi \supseteq \mathcal{P}(\mathbb{R})$. Let T be a transmutation operator and define $\varphi_k := T\left[x^k\right]$, $k \in \mathbb{N}_0$. From the transmutation property (5), we obtain

$$\left(\frac{d^2}{dx^2} - q\right)\varphi_k = \left(\frac{d^2}{dx^2} - q\right) Tx^k = T\frac{d^2}{dx^2}x^k.$$

Thus,

$$\left(\frac{d^2}{dx^2} - q\right)\varphi_k = 0, \quad k = 0, 1; \tag{6}$$

and

$$\left(\frac{d^2}{dx^2} - q\right)\varphi_k = k(k-1)\varphi_{k-2}, \quad k \geq 2. \tag{7}$$

As in [11], we define the following:

Definition 6 *A system of functions* $\{\varphi_k\}$, $k \in \mathbb{N}_0$, *satisfying (6) and (7) in the sense of weak solutions, is called an L-base.*

We have shown that if T is a transmutation operator on $\mathcal{P}(\mathbb{R})$, then $\varphi_k := T\left[x^k\right]$ is an L-base. Moreover, given an L-base $\{\varphi_k\}$, there exists a transmutation operator T_φ on $\mathcal{P}(\mathbb{R})$ such that $\varphi_k = T_\varphi\left[x^k\right]$ for all $k \in \mathbb{N}_0$. To see this, define T_φ on powers to be $T_\varphi\left[x^k\right] := \varphi_k$ and extend it to $\mathcal{P}(\mathbb{R})$ by linearity. It follows from Proposition 3 that each member of an L-base is a function from the space $W^{2,1}(a,b)$, hence T_φ maps $\mathcal{P}(\mathbb{R})$ into $W^{2,1}(a,b)$. Also, if $P = \sum_{k=0}^{m} \alpha_k x^k$, from (6) and (7) we get

$$(\frac{d^2}{dx^2} - q)T_\varphi P = \sum_{k=0}^{m} \alpha_k (\frac{d^2}{dx^2} - q)\varphi_k = \sum_{k=2}^{m} \alpha_k k(k-1)\varphi_{k-2} = T_\varphi P''.$$

This proves the existence of a transmutation operator on $\mathcal{P}(\mathbb{R})$ satisfying $T_\varphi\left[x^k\right] = \varphi_k$.

However, regarding possible applications, it is important to establish the existence of transmutations on larger domains than $\mathcal{P}(\mathbb{R})$, for example on $W^{2,1}(a,b)$, and investigate when they are bounded or invertible. Looking for the existence of such operators, we will work with a special class of L-bases, which we introduce below. We will assume from now on, without loss of generality, that $0 \in [a,b]$.

Definition 7 *A standard L-base (or, simply, an s-L-base) is an L-base* $\{\varphi_k\}$ *such that* $\varphi_k(0) = \varphi_k'(0) = 0$ *for all* $k \geq 2$. *A transmutation operator is called standard if* $\varphi_k := T\left[x^k\right]$ *is an s-L-base.*

Example 8 *The system of powers* $\left\{x^k\right\}$, $k \in \mathbb{N}_0$, *is an s-L-base corresponding to the operator* $L = \frac{d^2}{dx^2}$. *The shift operator* $Eu(x) = u(x - x_0)$, $x_0 \neq 0$ *is a transmutation operator (on* $W^{2,1}(\mathbb{R})$*) corresponding to* $L = \frac{d^2}{dx^2}$ *but it is not an s-transmutation operator.*

Remark 9 *We stress that an s-L-base* $\{\varphi_k\}$ *is completely determined by its first two elements* φ_0 *and* φ_1. *For instance, from Proposition 3 and by the variation of parameters formula, we see that the unique weak solution of (7) satisfying* $\varphi_k(0) = \varphi_k'(0) = 0$ *is given by*

$$\varphi_k(x) = k(k-1)\int_0^x G(x,s)\varphi_{k-2}(s)ds, \quad k \geq 2, \tag{8}$$

where

$$G(x,s) = \frac{\psi_0(s)\psi_1(x) - \psi_0(x)\psi_1(s)}{W(\psi_0,\psi_1)}, \tag{9}$$

$W(\psi_0,\psi_1) = \psi_0(x)\psi_1'(x) - \psi_0'(x)\psi_1(x)$, *and* ψ_0, ψ_1 *are any two linearly independent weak solutions of (6). Note that* $G(x,s)$ *does not depend on the choice of* ψ_0 *and* ψ_1.

Theorem 10 *Suppose* $\{\varphi_k\}$ *is an s-L-base and suppose* $T : L^1(a,b) \to L^1(a,b)$ *is a bounded linear operator such that* $T[x^k] = \varphi_k$, $k \in \mathbb{N}_0$. *Then* T *is an s-transmutation operator on* $W^{2,1}(a,b)$.

Proof. Let us prove that

$$T[u] = u(0)\varphi_0(x) + u'(0)\varphi_1(x) + \int_0^x G(x,s)T\left[u''\right](s)ds \tag{10}$$

for any $u \in W^{2,1}(a,b)$, where $G(x,s)$ is given by (9). Then (5) will follow from the application of $L = \frac{d^2}{dx^2} + q(x)$ to both sides of (10).

As T is linear and $T[x^k] = \varphi_k$ is an s-L-base, the application of T to $P = \sum_{k=0}^{m} \alpha_k x^k$ gives

$$\begin{aligned} T\left[P\right] &= \sum_{k=0}^{m} \alpha_k T_h[x^k] = \sum_{k=0}^{m} \alpha_k \varphi_k(x) \\ &= \alpha_0\varphi_0(x) + \alpha_1\varphi_1(x) + \int_0^x G(x,s) \sum_{k=2}^{m} k(k-1)\alpha_k \varphi_{k-2}(s)ds \\ &= P(0)\varphi_0(x) + P'(0)\varphi_1(x) + \int_0^x G(x,s)T\left[P''\right](s)ds, \end{aligned}$$

where we have used (8). According to Proposition 4, given $u \in W^{1,2}(a,b)$ there exists a sequence of polynomials P_n such that $P_n \to u$, $P_n' \to u'$ uniformly on $[a,b]$ and $P_n'' \to u''$ in $L_1(a,b)$, as $n \to \infty$. Then (10) follows from a simple limiting procedure using the continuity of T and the above relations,

$$\begin{aligned} T\left[u\right] &= \lim_{n\to\infty} T\left[P_n\right] = \lim_{n\to\infty} \left[P_n(0)\varphi_0(x) + P_n'(0)\varphi_0(x) + \int_0^x G(x,s)T\left[P_n''\right](s)ds\right] \\ &= u(0)\varphi_0(x) + u'(0)\varphi_1(x) + \int_0^x G(x,s)T\left[u''\right](s)ds. \end{aligned}$$

■

Remark 11 *It is easy to see that the above theorem is still true if we replace the boundedness of T on $L^1(a,b)$ by the boundedness from $W^{1,1}(a,b)$ to $L^1(a,b)$. However, in this case the transmutation property (5) holds on $W^{3,1}(a,b)$, instead of on $W^{2,1}(a,b)$.*

The following proposition establishes conditions under which the limit of a sequence of s-transmutation operators is an s-transmutation operator as well.

Proposition 12 *Let q_n be a convergent sequence in $L^1(a,b)$ with limit q. Let T be a bounded operator on $L^1(a,b)$ and T_n be a sequence of s-transmutation operators corresponding to $L_n = \frac{d^2}{dx^2} + q_n$. If for each $k = 0, 1, 2..$*

$$T_n\left[x^k\right] \to T\left[x^k\right] \text{ uniformly on } [a,b], \text{ as } n \to \infty,$$

then T is an s-transmutation operator on $W^{2,1}(a,b)$ corresponding to $L = \frac{d^2}{dx^2} + q$.

Proof. Write $\varphi_{k,n}(x) = T_n\left[x^k\right]$ and $\varphi_k(x) = T\left[x^k\right]$. To prove the theorem, it suffices to show that $\{\varphi_k\}$ is an s-L-base. Indeed, using the boundedness of T together with the mapping property $T\left[x^k\right] = \varphi_k(x)$, we conclude from Theorem 10 that T is an s-transmutation operator on $W^{2,1}(a,b)$.

Let us start by proving that $L[\varphi_0] = \varphi_0''(x) - q(x)\varphi_0(x) = 0$. Consider the functions $r_n(x)$ and $r(x)$ defined by

$$r_n(x) = \varphi_{0,n}(x) - \int_0^x (x-t)q_n(t)\varphi_{0,n}(t)dt, \tag{11}$$

$$r(x) = \varphi_0(x) - \int_0^x (x-t)q(t)\varphi_0(t)dt. \tag{12}$$

It is easy to see from (11) that $r''(x) = 0$ iff $L[\varphi_0] = 0$. In this way, as $L_n\left[\varphi_{0,n}\right] = \varphi_{0,n}''(x) - q_n(x)\varphi_{0,n}(x) = 0$ from (11), we get $r_n''(x) = 0$. Thus $r_n(x) = c_1 + c_2 x$, and taking into account the boundary values at $x = 0$, we see that

$$r_n(x) = \varphi_{0,n}(0) + \varphi_{0,n}^{'}(0)x. \tag{13}$$

Since, by assumption, $\varphi_{0,n}(x) = T_n[1] \to T[1] = \varphi_0(x)$ uniformly on $[a,b]$ as $n \to \infty$, we can see from (11) and (12) that $r_n(x) \to r(x)$ uniformly on $[a,b]$. From this and (13) we must have that $\varphi_{0,n}'(0) \to \varphi_0'(0)$ as $n \to \infty$, and $r(x) = \varphi_0(0) + \varphi_0'(0)x$. Thus $r''(x) = 0$ and from (12) we conclude that $L[\varphi_0] = 0$. In the same manner we can see that $L[\varphi_1] = 0$.

Let $G(x,s)$ be the function in (9) where ψ_0 and ψ_1 are linearly independent solutions of $L[\psi] = 0$ and denote by $G_n(x,s)$ the analogous function corresponding to the operator L_n. From the fact that $q_n \to q$ in $L^1(a,b)$ it is easy to see that $G_n(x,s) \to G(x,s)$ uniformly on $[a,b] \times [a,b]$ as $n \to \infty$. Hence, as $\varphi_{k,n}(x) \to \varphi_k(x)$ uniformly, as $n \to \infty$, for $k \geq 2$ we finally obtain

$$\begin{aligned} \varphi_k(x) &= \lim_{n\to\infty} \varphi_{k,n}(x) = k(k-1) \lim_{n\to\infty} \int_0^x G_n(x,s)\varphi_{k-2,n}(s)ds \\ &= k(k-1) \int_0^x G(x,s)\varphi_{k-2}(s)ds. \end{aligned}$$

This shows that $\{\varphi_k\}$ is an s-L-base. The theorem is proved. ■

4 New s-transmutations

In this section a method for constructing new s-transmutation operators on $\mathcal{P}(\mathbb{R})$ in terms of one single s-transmutation operator is presented.

Let us introduce the operators

$$P_{\pm}u(x) = \frac{u(x) \pm u(-x)}{2}, \quad Au(x) = \int_0^x u(s)ds, \tag{14}$$

mapping $\mathcal{P}(\mathbb{R})$ into itself. Note that the $P_{\pm}$ are pairwise projection operators, that is, $P_+^2 = P_+$, $P_-^2 = P_-$, $P_+ + P_- = I$, and $P_+P_- = P_-P_+ = 0$.

Proposition 13 *The relations* $\frac{d}{dx}P_{\pm} = P_{\mp}\frac{d}{dx}$ *and* $AP_{\pm} = P_{\mp}A$ *hold on* $\mathcal{P}(\mathbb{R})$.

Proof. Straightforward. ■

Proposition 14 *Let* T *be an s-transmutation operator on* $\mathcal{P}(\mathbb{R})$. *Then*

i) $T[1] = 0$ *if and only if* $T = TP_-$ *on* $\mathcal{P}(\mathbb{R})$;

ii) $T[x] = 0$ *if and only if* $T = TP_+$ *on* $\mathcal{P}(\mathbb{R})$;

iii) $T[1] = 0$ *and* $T[x] = 0$ *if and only if* $T \equiv 0$ *on* $\mathcal{P}(\mathbb{R})$.

Proof. We prove the direct assertions corresponding to *i*), *ii*) and *iii*). The reciprocal assertions are obvious. First, denote by $\varphi_k = T[x^k]$ the s-L-base corresponding to T. Let us prove *i*). Since $\varphi_0 = T[1] = 0$, it follows from Remark 9 that $\varphi_k = T[x^k] = 0$ for $k \geq 0$ even. Thus it is easy to see that $T[x^k] = TP_-[x^k]$ for all $k \geq 0$. From this and from the linearity of T and TP_- we conclude that $T[p(x)] = TP_-[p(x)]$ for every polynomial $p(x)$. Part *ii*) is proved by analogy. Part *iii*) is a consequence of *i*) and *iii*). For instance, since $T = TI = T(P_+ + P_-) = TP_+ + TP_- = 2T$, we have $T = 0$. ■

Corollary 15 *Let* T *and* M *be s-transmutation operators on* $\mathcal{P}(\mathbb{R})$ *such that* $T[1] = M[1]$ *and* $T[x] = M[x]$. *Then* $T = M$ *on* $\mathcal{P}(\mathbb{R})$.

Proof. This follows at once from part *iii*) of the above proposition. Setting $S = T - M$, we have that S is an s-transmutation operator and $S[1] = S[x] = 0$. Thus, $S \equiv 0$ on $\mathcal{P}(\mathbb{R})$. ■

Proposition 16 *Let* T *be an s-transmutation operator on* $\mathcal{P}(\mathbb{R})$. *Then:*

i) $T\frac{d}{dx}$ *is an s-transmutation operator on* $\mathcal{P}(\mathbb{R})$ *if and only if* $T[x] = 0$, *and in this case* $T\frac{d}{dx} = T\frac{d}{dx}P_-$ *on* $\mathcal{P}(\mathbb{R})$;

ii) TA *is an s-transmutation operator on* $\mathcal{P}(\mathbb{R})$ *if and only if* $T[1] = 0$, *and in this case* $TA = TAP_+$ *on* $\mathcal{P}(\mathbb{R})$;

Proof. Let T be an s-transmutation operator and denote by $\varphi_k = T[x^k]$ the corresponding s-L-base. Let us prove *i*). The transmutation property of $T\frac{d}{dx}$ is a straightforward calculation. Analyzing the corresponding L-base $\psi_k = T\frac{d}{dx}[x^k] = kT\left[x^{k-1}\right] = k\varphi_{k-1}$, it is easy to conclude that ψ_k is standard iff $\varphi_1 = T[x] = 0$. Thus, from Proposition 14, $T = TP_+$ and therefore $T\frac{d}{dx} = TP_+\frac{d}{dx} = T\frac{d}{dx}P_-$.

(*ii*) From the equalities $(\frac{d^2}{dx^2} - q)TAu = T\frac{d^2}{dx^2}Au = T\frac{d}{dx}u$ and

$$TA\tfrac{d^2}{dx^2}u(x) = T\left[u'(x) - u'(0)\right] = T\tfrac{d}{dx}u - u'(0)T[1]$$

we see that TA is a transmutation operator on $\mathcal{P}(\mathbb{R})$ whenever $T[1] = 0$. In this case, Proposition 14 tell us that $T = TP_+$, from which follows the equality $TA = TP_+A = TAP_-$. ■

Theorem 17 *Let T be an s-transmutation operator on $\mathcal{P}(\mathbb{R})$. Then the operators TP_+, TP_-, TAP_+ and $T\frac{d}{dx}P_-$ are s-transmutations on $\mathcal{P}(\mathbb{R})$ as well.*

Proof. Using the relation $\frac{d^2}{dx^2}P_+ = P_+\frac{d^2}{dx^2}$ (Proposition 13) it is easy to see that TP_+ is a transmutation operator. As the corresponding L-base $\psi_k = TP_+[x^k]$ is such that $\psi_k = T[x^k] = \varphi_k$ for k even, and $\psi_k = 0$ for k odd, we easily see that $\psi_k(0) = \psi_k'(0) = 0$, $k \geq 2$. Thus ψ_k is an s-L-base and TP_+ is an s-transmutation operator.

Let us now consider the operator $T\frac{d}{dx}P_-$. As TP_+ is an s-transmutation and $TP_+[x] = 0$, then, in accordance with Proposition 16 part i), the operator $TP_+\frac{d}{dx} = T\frac{d}{dx}P_-$ is an s-transmutation as well.

By analogous means we can prove that TP_- and TAP_+ are s-transmutation operators. ■

Proposition 18 *Let $\{\varphi_k\}$ and $\{\psi_k\}$ be s-L-bases such that $W(\psi_0, \psi_1) \neq 0$, and let T_φ, T_ψ be the corresponding s-transmutation operators satisfying $T_\varphi[x^k] = \varphi_k$, $T_\psi[x^k] = \psi_k$. Then*

$$T_\varphi = \tfrac{W(\varphi_0,\psi_1)}{W(\psi_0,\psi_1)}T_\psi P_+ - \tfrac{W(\varphi_0,\psi_0)}{W(\psi_0,\psi_1)}T_\psi AP_+ + \tfrac{W(\varphi_1,\psi_1)}{W(\psi_0,\psi_1)}T_\psi \tfrac{d}{dx}P_- - \tfrac{W(\varphi_1,\psi_0)}{W(\psi_0,\psi_1)}T_\psi P_- \qquad (15)$$

where the equality is valid on $\mathcal{P}(\mathbb{R})$.

Proof. Let T denote the operator defined on $\mathcal{P}(\mathbb{R})$ by the right hand side of (15). Note that as $\varphi_0, \varphi_1, \psi_0$ and ψ_1 are solutions of $u'' = qu$, all the involved Wronskians on (15) $W(u_i, v_i)$ are constant functions and thus T is an s-transmutation operator on $\mathcal{P}(\mathbb{R})$ since it is a linear combination of the s-transmutations $T_\psi P_+$, $T_\psi AP_+$, $T_\psi \frac{d}{dx}P_-$ and $T_\psi P_-$ (see Theorem 17). As $T_\psi P_-[1] = T_\psi \frac{d}{dx}P_-[1] = T_\psi P_+[x] = T_\psi AP_+[x] = 0$, it follows that

$$T[1] = \tfrac{W(\varphi_0,\psi_1)}{W(\psi_0,\psi_1)}\psi_0 - \tfrac{W(\varphi_0,\psi_0)}{W(\psi_0,\psi_1)}\psi_1 = \varphi_0 = T_\varphi[1]$$

and

$$T[x] = \tfrac{W(\varphi_1,\psi_1)}{W(\psi_0,\psi_1)}\psi_0 - \tfrac{W(\varphi_1,\psi_0)}{W(\psi_0,\psi_1)}\psi_1 = \varphi_1 = T_\varphi[x].$$

Lastly, as T_φ and T are s-transmutations and $T[1] = T_\varphi[1]$, $T[x] = T_\varphi[x]$, it follows from Corollary 15 that $T_\varphi = T$ on $\mathcal{P}(\mathbb{R})$. ■

5 s-Transmutation operators on Sobolev spaces

5.1 Existence of s-transmutation operators on $\mathbf{W}^{2,1}(-a, a)$

Let $q \in C[-a, a]$, $a > 0$. Then, according to [27], a transmutation operator on $C^2[-a, a]$ corresponding to $L = \frac{d^2}{dx^2} - q(x)$ can be represented in the form

$$\mathbf{T}u(x) = u(x) + \int_{-x}^{x} K(x, t)u(t)dt$$

where $K(x,t)$ is constructed as follows. Consider the transformation $2u = x+t$, $2v = x-t$, which maps the square $\Omega : -a \leq x, y \leq a$ onto the square Ω_1 in the (u,v)-plane with vertices $(-a,0),(0,a),(a,0)$ and $(0,-a)$. Then, $K(x,t) = H(\frac{x+t}{2}, \frac{x-t}{2}) = H(u,v)$ where $H(u,v)$ is the unique solution of the integral equation

$$H(u,v) = \frac{1}{2}\int_0^u q(s)ds + \int_0^u \int_0^v q(\alpha+\beta)H(\alpha,\beta)d\beta d\alpha, \text{ in } \Omega_1. \tag{16}$$

If q is n times differentiable, then $K(x,t)$ is $n+1$ times continuously differentiable [27]. Moreover, if $q \in C^1[-a,a]$, then $K(x,t) \in C^2(\Omega)$ is the unique classical solution of the following Goursat problem

$$\left(\frac{\partial^2}{\partial x^2} - q(x)\right) K(x,t) = \frac{\partial^2}{\partial t^2} K(x,t), \quad \text{in } \Omega. \tag{17}$$

$$K(x,x) = \frac{1}{2}\int_0^x q(s)ds, \;\; K(x,-x) = 0. \tag{18}$$

We stress that if q is differentiable, then the transmutation property of $\mathbf{T}$ follows from straightforward calculations, using (17) and (18), [23]. However, for a continuous q, since $K(x,t) \notin C^2(\Omega_1)$, the proof of this result is not trivial. In [27], a proof involving the use of Riemann's function was presented. Another method was mentioned in [25, pag.9] without presenting a rigorous proof: if we approximate the continuous potential q by differentiable potentials q_n, then the transmutation property of $\mathbf{T}$ should follow by a limit extrapolation of the transmutations $\mathbf{T}_n$, say $\mathbf{T} = \lim \mathbf{T}_n$. This idea was also used in [21, proof of Th. 3.4] and [22, proof of Th. 6].

In this section, we give a rigorous justification of the procedure proposed in [25], and combining it with Proposition 12, we prove the existence of a transmutation operator on $W^{2,1}(-a,a)$, in the case when $q \in L^1(-a,a)$.

Proposition 19 *Let $q \in L^1(-a,a)$. Then the integral equation (16) has a unique solution $H(u,v)$. Moreover, $H(u,v) \in C(\Omega_1) \cap W^{1,1}(\Omega_1)$ and*

$$|H(u,v)| \leq \|q\|_{L^1} e^{a\|q\|_{L^1}}. \tag{19}$$

Proof. The proof is analogous to the one given in [27, Th. 1.2.2] for a continuous potential q. For convenience, we sketch the main steps. First, using the method of successive approximations, we write a hypothetical series representation for the solution of (16), say $H(u,v) = \sum_{n=0}^{+\infty} H_n(u,v)$ where $H_0(u,v) = \frac{1}{2}\int_0^u q(s)ds$ and $n \geq 1$

$$H_n(u,v) = \int_0^u \int_0^v q(\alpha+\beta)H_{n-1}(\alpha,\beta)d\beta d\alpha. \tag{20}$$

Next, we prove the uniform convergence of the above series, which in turn will imply the existence and the unicity of a continuous solution of (16). From (20), and following [27], we arrive at the following estimate

$$|H_n(u,v)| \leq \|H_0\|_{C(\Omega_1)} \frac{1}{n_!} \left| \int_0^{u+v} \left| \int_0^\beta |q(\alpha)|\, d\alpha \right| d\beta \right|^n \leq \|q\|_{L^1} \frac{(a\|q\|_{L^1})^n}{n!}$$

From this and by the Weierstrass M-test it follows that the series defining H is uniformly convergent and $|H(u,v)| \leq \sum_{n=0}^{+\infty} |H_n(u,v)| \leq \|q\|_{L^1} e^{a\|q\|_{L^1}}$. Lastly, from (16) we see that $H \in W^{1,1}(\Omega_1)$ as it is the sum of the absolutely continuous function $\frac{1}{2}\int_0^u q(s)ds$ with a function in $C^1(\Omega_1)$. ■

Proposition 20 *Let $q \in L^1(-a,a)$ and let $H(u,v)$ be the unique solution of (16). Then the mapping $q \to H(u,v)$ is continuous from $L^1(-a,a)$ to $C(\Omega_1)$.*

Proof. Denote by $\widetilde{H}(u,v)$ the solution of (16) with q replaced by $\widetilde{q} \in L^1(-a,a)$. It suffices to prove that the inequality

$$\left|\widetilde{H}(u,v) - H(u,v)\right| \leq C \|\widetilde{q} - q\|_{L^1} e^{a\|\widetilde{q}\|_{L^1}}$$

holds for all $q, \widetilde{q} \in L^1(-a,a)$ with some constant $C > 0$ not depending on $\widetilde{q}$. In order to prove this, we first observe that $M(u,v) := \widetilde{H}(u,v) - H(u,v)$ is the unique solution of the integral equation $M(u,v) = M_0(u,v) + \int_0^u \int_0^v \widetilde{q}(\alpha+\beta)M(\alpha,\beta)d\beta d\alpha$ where

$$M_0(u,v) = \frac{1}{2}\int_0^u (\widetilde{q}(s) - q(s))\, ds + \int_0^u \int_0^v H(\alpha,\beta)\,(\widetilde{q}(\alpha+\beta) - q(\alpha+\beta))\, d\beta d\alpha. \tag{21}$$

Now, applying the same reasoning as in the proof of the above proposition, one can check that $|M(u,v)| \leq \|M_0\|_{C(\Omega_1)} e^{a\|\widetilde{q}\|_{L^1}}$. On the other hand, from (21) we get

$$|M_0(u,v)| \leq \frac{1}{2}\|\widetilde{q} - q\|_{L^1} + a\sup_{\Omega_1} |H(\alpha,\beta)|\, \|\widetilde{q} - q\|_{L^1} = C\|\widetilde{q} - q\|_{L^1}.$$

Now, combining the last two inequalities, the desired result follows. ■

Theorem 21 *Let $q \in L^1(-a,a)$ and let $H(u,v)$ be the unique solution of (16). Then, $K(x,t) = H(\frac{x+t}{2}, \frac{x-t}{2})$ is a weak solution of the Goursat problem (17), (18) and*

$$\mathbf{T}u(x) = u(x) + \int_{-x}^{x} K(x,t)u(t)dt \tag{22}$$

is an s-transmutation operator on $W^{2,1}(-a,a)$, corresponding to $L = \frac{d^2}{dx^2} - q(x)$. The corresponding L-base $\alpha_k(x) := \mathbf{T}\left[x^k\right]$ is such that

$$\alpha_0(0) = \alpha_1'(0) = 1 \text{ and } \alpha_0'(0) = \alpha_1(0) = 0. \tag{23}$$

Moreover, $\mathbf{T}$ is invertible and $\mathbf{T}^{-1}u(x) = u(x) - \int_{-x}^{x} K(t,x)u(t)dt$.

Proof. Let $q_n \in C^1[-a,a]$ be a sequence of functions that converges to q in $L^1(-a,a)$ and let $H_n(u,v)$ be the unique solution of the integral equation

$$H_n(u,v) = \frac{1}{2}\int_0^u q_n(s)ds + \int_0^u \int_0^v q_n(\alpha+\beta)H_n(\alpha,\beta)d\beta d\alpha. \tag{24}$$

Since $q_n \in C^1[-a,a]$, we have that $K_n(x,t) := H_n(\frac{x+t}{2}, \frac{x-t}{2})$ is a classical a solution of the equation $(\partial_x^2 - \partial_t^2 - q_n(x))K_n(x,t) = 0$ (see the beginning of this section). In particular, the equality

$$\int\int_\Omega \left[(\partial_t K_n)\,\partial_t\varphi - (\partial_x K_n)\,\partial_x\varphi - q_n K_n\varphi\right] dxdt = 0, \tag{25}$$

holds for all $\varphi \in C_c^\infty(\Omega)$. As $q_n \to q$ in $L^1(-a,a)$, it follows from Proposition 20 that $H_n \to H$ uniformly on Ω_1. Moreover, differentiating (24) and using the uniform convergence of H_n, we see that $\partial_u H_n \to \partial_u H$ and $\partial_v H_n \to \partial_v H$ in $L^1(\Omega_1)$ as $n \to \infty$. Thus $K_n \to K$ uniformly on Ω and $\partial_x K_n \to \partial_x K$, $\partial_y K_n \to \partial_y K$ in $L^1(\Omega)$ as $n \to \infty$. Then, letting $n \to +\infty$ in (25), we obtain

$$\int\int_\Omega \left[(\partial_t K)\,\partial_t\varphi - (\partial_x K)\,\partial_x\varphi - qK\varphi\right] dxdt = 0$$

for all $\varphi \in C_c^\infty(\Omega)$. Thus K is a weak solution of (17). The boundary conditions (18) for K are obtained by taking the limit of the corresponding boundary conditions for K_n, $K_n(x,x) = \frac{1}{2}\int_0^x q_n(s)ds$, $K_n(x-x) = 0$. Thus K is a weak solution of (17), (18).

The operator $\mathbf{T}$ is an s-transmutation because it is the uniform limit of the s-transmutation operators $\mathbf{T}_n u(x) = u(x) + \int_{-x}^x K_n(x,t)u(t)dt$, see Proposition 12. The initial conditions (23) of the L-base $\{\alpha_k\}$ follow by differentiating (22) and then using (18) at $x = 0$.

The formula for the kernel of T^{-1} was obtained in [22] for a continuous potential q and, by using a limit procedure as above, it is easy to see that it remains true for an integrable q. The theorem is proved. ■

5.2 A fundamental set of s-transmutation operators

Let $\mathbf{T}$ be the s-transmutation operator (22) studied in Proposition 21. In accordance with Theorem 17 each one of the operators

$$\mathbf{T}P_+,\ \mathbf{T}P_-,\ \mathbf{T}AP_+,\ \mathbf{T}\tfrac{d}{dx}P_-, \tag{26}$$

is an s-transmutation on $\mathcal{P}(\mathbb{R})$ (the linear space of polynomials). Next we show that (26) is a fundamental set of s-transmutation operators in the sense that any s-transmutation operator is a linear combination of operators from (26). Set $I = (-a,a)$, $0 < a < \infty$.

Theorem 22 *Let $\{\varphi_k\}$, $k \in \mathbb{N}_0$, be an s-L-base. Then there exists T_φ an s-transmutation operator on $W^{3,1}(I)$ such that $T_\varphi[x^k] = \varphi_k$. The equality*

$$T_\varphi = \varphi_0(0)\mathbf{T}P_+ + \varphi_0'(0)\mathbf{T}AP_+ + \varphi_1(0)\mathbf{T}\tfrac{d}{dx}P_- + \varphi_1'(0)\mathbf{T}P_- \tag{27}$$

holds on $W^{1,1}(I)$, where the operators $\mathbf{T}$, $P_{\pm}$, and A are given by (22) and (14). Also, $T_\varphi : W^{1,1}(I) \to L^1(I)$ is bounded. Moreover, if $W(\varphi_0, \varphi_1) \neq 0$, then the operator

$$M_\varphi := \frac{1}{W(\varphi_0, \varphi_1)} \left\{ \varphi_1'(0) P_+ \mathbf{T}^{-1} - \varphi_0'(0) A P_+ \mathbf{T}^{-1} - \varphi_1(0) \tfrac{d}{dx} P_- \mathbf{T}^{-1} + \varphi_0(0) P_- \mathbf{T}^{-1} \right\}$$

satisfies $T_\varphi M_\varphi(u) = M_\varphi T_\varphi(u) = u$ for all $u \in W^{2,1}(I)$. In particular, if $\varphi_1(0) = 0$, then $T_\varphi : L^1(I) \to L^1(I)$ is bounded, and if in addition $\varphi_0(0) \neq 0$, then T_φ is invertible with $T_\varphi^{-1} = M_\varphi$.

Proof. As $\mathbf{T}$ and T_φ are both s-transmutations on $\mathcal{P}(\mathbb{R})$ with $T_\varphi[x^k] = \varphi_k$ and $\mathbf{T}[x^k] = \alpha_k$, from Proposition 18 and using (23) we see that (27) holds on $\mathcal{P}(\mathbb{R})$. As $\mathbf{T} : L^1(I) \to L^1(I)$ is bounded, the operator defined on $\mathcal{P}(\mathbb{R})$ by the right hand side of (27) can be extended continuously to $W^{1,1}(I)$, from which we conclude that $T_\varphi : W^{1,1}(I) \to L^1(I)$ is bounded. Thus employing Theorem 10 and Remark 11, T_φ is an s-transmutation operator on $W^{3,1}(I)$. The relation $T_\varphi M_\varphi(u) = M_\varphi T_\varphi(u) = u$ is obtained after a few but trivial calculations, and using the existence of $\mathbf{T}^{-1}$ (see Proposition 21). All the remaining affirmations are obvious. ■

Corollary 23 *Let $\{\varphi_k\}$ be an s-L-base such that $W(\varphi_0, \varphi_1) \neq 0$ and $\lambda \in \mathbb{C}$. Then the general weak solution of the equation*

$$v'' + qv = \lambda v \tag{28}$$

on I has the form $v_{ger} = c_1 v_1 + c_2 v_2$ where c_1 and c_2 are arbitrary complex constants,

$$v_1 = \sum_{k=0}^{\infty} \lambda^k \frac{\varphi_{2k}}{(2k)!} \quad \text{and} \quad v_2 = \sum_{k=1}^{\infty} \lambda^k \frac{\varphi_{2k+1}}{(2k+1)!}, \tag{29}$$

and both series, as well as the corresponding series of derivatives, converge uniformly on $\overline{I}$.

Proof. Let T_φ be the s-transmutation operator studied in Theorem 22 and satisfying $T_\varphi[x^k] = \varphi_k$. From the transmutation property of T_φ, we can see that $(\frac{d^2}{dx^2} - q - \lambda) T_\varphi u = T_\varphi(\frac{d^2}{dx^2} - \lambda) u$ for all $\lambda \in \mathbb{C}$ and $u \in W^{3,1}(I)$. Thus, the operator T_φ maps classical solutions of $u'' = \lambda u$ to weak solutions of (28). Moreover, as $W(\varphi_0, \varphi_1) \neq 0$, it follows that T_φ is one-to-one on $W^{3,1}(I)$, and thus the general solution of (28) is obtained by applying T_φ to the general solution of $u'' = \lambda u$, $u_{ger} = c_1 \cosh(\lambda x) + c_2 (1/\sqrt{\lambda}) \sinh(\sqrt{\lambda} x)$. Thus, $v_{ger} = T_\varphi[u_{ger}] = c_1 T_\varphi[\cosh(\sqrt{\lambda} x)] + c_2 T_\varphi[(1/\sqrt{\lambda}) \sinh(\sqrt{\lambda} x)]$ where

$$T_\varphi[\cosh(\lambda x)] = T_\varphi \left[\sum_{k=0}^{+\infty} \lambda^k \frac{x^{2k}}{(2k)!} \right] = \sum_{k=0}^{+\infty} \lambda^k \frac{\varphi_{2k}}{(2k)!} = v_1$$

and $T_\varphi[(1/\sqrt{\lambda}) \sinh(\sqrt{\lambda} x)] = T_\varphi \left[\sum_{k=0}^{+\infty} \lambda^k \frac{x^{2k+1}}{(2k+1)!} \right] = \sum_{k=0}^{+\infty} \lambda^k \frac{\varphi_{2k+1}}{(2k+1)!} = v_2$. Lastly, the convergence of the series defining the solutions v_1 and v_2 follows from the fact that T_φ is bounded from $C^2(\overline{I})$ into $C^1(\overline{I})$. ■

Remark 24 *Let us relate the results we have obtained so far with the ones appearing in the literature. Assume that* $q \in C\left(\overline{I}\right)$ *and that equation (6) possesses a nonvanishing solution* $f \in C^2(\overline{I})$. *Set* $\varphi_0 := f$ *and note that a second linearly independent solution can be computed by the formula* $\varphi_1(x) = f(x)\int_0^x \frac{dt}{f^2(t)}$. *Without loss of generality assume that* $f(0) = 1$ *and set* $c := f'(0)$. *Thus,* $\varphi_0(0) = 1$, $\varphi_0'(0) = c$, $\varphi_1(0) = 0$, *and* $\varphi_1'(0) = 1$. *Let* $\{\varphi_k\}$ *be the s-L-base corresponding to this pair of solutions, and denote by* T_c *the (transmutation) operator such that* $T_c\left[x^k\right] = \varphi_k$. *Notice that* $T_0 \equiv \mathbf{T}$. *The operator* T_c *has been studied in several places in the literature. In [8], assuming* $q \in C^1(\overline{I})$, *it was proved that* T_c *is a Volterra integral operator and also a transmutation operator on* $C^2(\overline{I})$. *Another proof of the transmutation property of* T_c, *for any value of* $c \in \mathbb{C}$, *was given in [22] when assuming that* $q \in C^1(\overline{I})$ *and also on [23] supposing that* $q \in C(\overline{I})$ *and that* $\varphi_0(x)$ *is nonvanishing on* $\overline{I}$. *In [8] a formula for the kernel of the operator* T_c *was obtained in terms of the kernel of the same operator with* $c = 0$. *Starting from this result, the relation* $T_c u = \mathbf{T}\left[u(x) + \frac{c}{2}\int_{-x}^{x} u(t)dt\right]$ *was obtained in [23]. This equality is the one given by (27), in this special case. In [21], and with the help of the above relation, it was proved that* T_c *is a transmutation operator for any* $q \in C(\overline{I})$ *and for any value of* c. *Perhaps at the same time, another proof for this result was presented in [6], which is published here (the proof of Theorem 10) for the first time.*

The spectral parameter power series representation, like those representing the general solution of (28) in Corollary 23, was first introduced in [17] as an application of pseudoanalytic function theory and nowadays represents a useful tool for the numerical solution of both regular and singular Sturm–Liouville problems, see [4], [21] and the references therein. Originally, the coefficients of the power series (29) were computed by slightly different recursive formulas, and its construction required the existence of a nonvanishing solution of (28) with $\lambda = 0$. *Here we have relaxed this restriction using the s-L-bases framework.*

6 s-Transmutation operators on distribution spaces

Let $I = (-a, a)$, $0 < a < \infty$, and let $q \in L^1_{loc}(I)$. A linear operator $T : D_T \subset \mathcal{D}'(I) \to \mathcal{D}'(I)$ is called a transmutation operator on $\chi \subset \mathcal{D}'(I)$ if for every $u \in \chi$

$$\left(\frac{d^2}{dx^2} - q\right) Tu = T\frac{d^2}{dx^2}u \quad \text{in } \mathcal{D}'(I).$$

Our next goal is to show that the transmutation operators constructed in Section 5 can be continuously extended from Sobolev spaces to suitable spaces of distributions, preserving the transmutation property. To this, let us consider the s-transmutation operator studied in Section 5.1, Theorem 21,

$$\mathbf{T}u(x) = u(x) + \int_{-x}^{x} K(x,t)u(t)dt. \tag{30}$$

Since $q \in L^1_{loc}(I)$, we have $K(x,t) \in C(I \times I) \cap W^{1,1}_{loc}(I \times I)$ and thus $\mathbf{T}$ maps $L^1_{loc}(I)$ into itself. In order to extend $\mathbf{T}$ to distributions, using Proposition 2, we first have to look for

the existence of the corresponding transpose operator. An easy computation shows that

$$\mathbf{T}^{\square}\psi(x) = \psi(x) - \int_{-a}^{-|x|} K(t,x)\psi(t)dt + \int_{|x|}^{a} K(t,x)\psi(t)dt, \quad \psi \in C_c(I), \tag{31}$$

with the equality $\int_I (\mathbf{T}u)\phi dx = \int_I u(\mathbf{T}^{\square}\phi)dx$ valid for all $u \in L^1_{loc}(I)$, $\phi \in C_c(I)$.

Let us treat first the case when $q \in C^\infty(I)$.

Proposition 25 *Suppose $q \in C^\infty(I)$. If the operator T is continuous on $\mathcal{D}'(I)$, and a transmutation on $C_c^\infty(I)$, then T is a transmutation operator on $\mathcal{D}'(I)$.*

Proof. Let $u \in \mathcal{D}'(I)$. As $C_c^\infty(I)$ is dense in $\mathcal{D}'(I)$ [12, Th. 3.18], there exists a sequence $\phi_n \in C_c^\infty(I)$ such that $\phi_n \to u$ in $\mathcal{D}'(I)$ as $n \to \infty$. Hence,

$$\left(\frac{d^2}{dx^2} - q\right)\mathbf{T}u = \lim_{n\to\infty}\left(\frac{d^2}{dx^2} - q\right)\mathbf{T}p_n = \lim_{n\to\infty}\mathbf{T}\frac{d^2}{dx^2}p_n = \mathbf{T}\frac{d^2}{dx^2}u, \quad \text{in } \mathcal{D}'(I),$$

where we have used the continuity in $\mathcal{D}'(I)$ of the involved operators and the transmutation property of T on $C_c^\infty(I)$. ■

Proposition 26 *Assume that $q \in C^\infty(I)$ and let $\mathbf{T}$ be the linear operator acting on distributions according to the rule*

$$\langle \mathbf{T}u, \phi\rangle := \left\langle u, \mathbf{T}^{\square}\phi\right\rangle, \quad u \in \mathcal{D}'(I), \phi \in C_c^\infty(I) \tag{32}$$

where $\mathbf{T}^{\square}$ is given by (31). Then, $\mathbf{T} : \mathcal{D}'(I) \to \mathcal{D}'(I)$ is continuous and invertible. Also, $\mathbf{T}$ is a transmutation operator on $\mathcal{D}'(I)$.

Proof. First let us prove that $\mathbf{T}$ is well defined and continuous on $\mathcal{D}'(I)$. According to Proposition 2, it suffices to show that $\mathbf{T}^{\square}$ is continuous on $C_c^\infty(I)$. Suppose the sequence ϕ_n converge in $C_c^\infty(I)$ to ϕ. Then there exists a number $b < a$ such that $\operatorname{supp}\phi_n \subseteq [-b,b] \subset I$ for all n, and from (31) we see that $\operatorname{supp}(\mathbf{T}^{\square}\phi_n) \subseteq [-b,b] \subset I$ for all n. On the other hand, as $q \in C^\infty(I)$, we have $K(x,t) \in C^\infty(I \times I)$ and thus the expression (31) defining $\mathbf{T}^{\square}\phi$ can be differentiated infinitely many times whenever $\phi \in C_c^\infty(I)$. For example,

$$\begin{aligned}\frac{d}{dx}\mathbf{T}^{\square}\phi(x) = \frac{d}{dx}\phi(x) - \int_{-a}^{-|x|}\partial_x K(t,x)\phi(t)dt + \int_{|x|}^{a}\partial_x K(t,x)\phi(t)dt \\ + K(-x,x)\phi(-x) + K(x,x)\phi(x).\end{aligned} \tag{33}$$

From the above facts it is easy to see that $\mathbf{T}^{\square}$ maps $C_c^\infty(I)$ into itself and that $\mathbf{T}^{\square}\phi_n \to \mathbf{T}^{\square}\phi$ in $C_c^\infty(I)$ whenever $\phi_n \to \phi$ in $C_c^\infty(I)$, as $n \to \infty$. This proves the continuity of $\mathbf{T}^{\square}$.

Since $\mathbf{T}$ is invertible on $L^1_{loc}(I)$ and $\mathbf{T}^{-1}$ is a Volterra integral operator (Theorem 21), it follows that $\mathbf{T}^{\square}$ is invertible as well, and $\left(\mathbf{T}^{\square}\right)^{-1} = (\mathbf{T}^{-1})^{\square}$. Therefore the inverse operator of $\mathbf{T}$ on $\mathcal{D}'(I)$ is defined by the rule $\langle \mathbf{T}^{-1}u, \phi\rangle := \left\langle u, \left(\mathbf{T}^{\square}\right)^{-1}\phi\right\rangle$ for $u \in \mathcal{D}'(I)$ and $\phi \in C^\infty(I)$.

Lastly, the transmutation property of $\mathbf{T}$ on $\mathcal{D}'(I)$ follows from Proposition 25 upon observing that $\mathbf{T}$ is continuous on $\mathcal{D}'(I)$ and a transmutation on $C_c^\infty(I)$ (Theorem 21). ■

Corollary 27 *Let $q \in C^\infty(I)$. Then the differential equation $u'' = qu$, $u \in \mathcal{D}'(I)$ has only classical solutions.*

Proof. Let $u \in \mathcal{D}'(I)$ be a solution of $u'' = qu$. Then, from the transmutation property of $\mathbf{T}$ it follows that $v = \mathbf{T}^{-1}u$ is a solution of $v'' = 0$, $v \in \mathcal{D}'(I)$. The last equation has only classical solutions [2], thus $v = C_1 + C_2 x$ for some constants C_1, C_2 and therefore $u = \mathbf{T}v = C_1\mathbf{T}[1] + C_2\mathbf{T}[x] \in C^\infty(I)$ is a classical solution of the first equation. ■

Corollary 28 *Let $q \in C^\infty(I)$. If $\mathcal{T}$ is a transmutation operator on $\mathcal{D}'(I)$, then $\mathcal{T}\left[x^k\right]$, $k \in \mathbb{N}_0$, is a regular distribution from the space $C^\infty(I)$.*

Proof. Set $\varphi_k := \mathcal{T}\left[x^k\right] \in \mathcal{D}'(I)$ for $k \in \mathbb{N}_0$. From the transmutation property of $\mathcal{T}$ it follows that φ_k is a distributional solution of (6) when $k = 0, 1$. For $k \geq 2$, it satisfies (7). However, from Corollary 27 we see that Equations (6) and (7) have only classical solutions, thus $\mathcal{T}\left[x^k\right] \in C^\infty(I)$. ■

The last result allows one to consider the notion of standard transmutation operator (Definition 7) on spaces of distributions.

In what follows, we obtain the analogue of some results of Section 4 for distributions. Consider the operators $P_\pm u(x) = \frac{u(x) \pm u(x)}{2}$ and $Au(x) = \int_0^x u(t)dt$ acting on $L^1_{loc}(I)$. Since $P_\pm^\square = P_\pm$ is continuous on $C_c^\infty(I)$, it follows from Proposition 2 that $P_\pm$ is well defined and continuous on $\mathcal{D}'(I)$. A similar extension is not available for the operator A. Indeed,

$$A^\square\psi(x) = \begin{cases} -\int_{-a}^x \psi(t)dt & , -a \leq x \leq 0 \\ \int_x^a \psi(t)dt & 0 < x \leq a \end{cases}.$$

Thus if $\psi \in C_c^\infty(I)$, we can not be assured of the continuity of the function $A^\square\psi(x)$ at $x = 0$. However, it is easy to see that $(AP_+)^\square = (P_-A)^\square = A^\square P_-^\square = A^\square P_-$ is continuous on $C_c^\infty(I)$, and thus the operator AP_+ admits a continuous extension to $\mathcal{D}'(I)$.

Proposition 29 *Let $q \in C^\infty(I)$ and let $\mathbf{T}$ be the operator defined in $\mathcal{D}'(I)$ by (30)-(32). Then, a continuous linear operator $\mathcal{T} : \mathcal{D}'(I) \to \mathcal{D}'(I)$ is an s-transmutation on $\mathcal{D}'(I)$ if and only if $\mathcal{T}$ is a linear combination of operators from (26).*

Proof. Once we have shown that each operator in (26) is a continuous s-transmutation on $\mathcal{D}'(I)$, then the same will be true for all their linear combinations. The continuity follows from the fact that each one of the operators $\mathbf{T}$, $P_\pm$, $\frac{d}{dx}$, and AP_+ is continuous on $\mathcal{D}'(I)$, and the transmutation property on $\mathcal{D}'(I)$ is a consequence of Theorem 22 and Proposition 25.

Conversely, let $\mathcal{T}$ be a continuous s-transmutation operator on $\mathcal{D}'(I)$. From Corollary 28 we see that the corresponding L-base, $\varphi_k := \mathcal{T}\left[x^k\right]$ is such that $\varphi_k \in C^\infty(I)$. Let us set

$$T_\varphi := \varphi_0(0)\mathbf{T}P_+ + \varphi_0'(0)\mathbf{T}AP_+ + \varphi_1(0)\mathbf{T}\tfrac{d}{dx}P_- + \varphi_1'(0)\mathbf{T}P_-.$$

Note that by Theorem 22, we have $T_\varphi\left[x^k\right] = \varphi_k$. Thus, $T_\varphi\left[x^k\right] = \mathcal{T}\left[x^k\right]$ for every $k \in \mathbb{N}_0$, and as both operators are linear they are equal on polynomials. In fact, as the set of

polynomials is dense in $\mathcal{D}'(I)$ and both operators are continuous on $\mathcal{D}'(I)$, we see that T_φ and $\mathcal{T}$ are equal on $\mathcal{D}'(I)$. This shows that $\mathcal{T} \equiv T_\varphi$ is a linear combination of operators from (26). ■

Now assume that $q \notin C^\infty(I)$. In such a case, the domain of the operator $\mathbf{T}$ defined in (30) is smaller than $\mathcal{D}'(I)$ and it depends on the smoothness of the kernel $K(x,t)$, which in turn depends on the smoothness of the potential q. Denote by $F(I)$, $I = (-a,a)$, any one of the following spaces $L^p(I)$, $C^k(I)$, $k \in \mathbb{N}_0$, $1 \leq p \leq \infty$. We are going to consider $F(I)$ with the convergence induced by the inductive limit topology (see Section 2). Define $W^2(I;F) = \{\phi \in W^{1,1}(I) : \phi'' \in F(I)\}$. If $K \subset I$ is compact, then $W^2(K;F)$ is a Banach space with norm $\|u\|_{W^2(K,F)} = \|u\|_{W^{1,1}(K)} + \|u\|_{F(k)}$. We shall also consider $W_c^2(I;F)$ with the inductive limit topology.

The extension of an operator T from functions to functionals will be based once more on the identity $\langle Tu, \phi\rangle = \left\langle u, T^\square \phi\right\rangle$, whenever it makes sense.

Proposition 30 *Assume that $q \in F_{loc}(I)$. Then the operator $\mathbf{T}$ defined in (30) is continuous and invertible on $(W_c^2(I;F))'$ as well on $(F_c(I))'$. Moreover, $\mathbf{T}$ is a transmutation operator on $(F_c(I))'$.*

Proof. The continuity of $\mathbf{T}$ on $(W_c^2(I;F))'$ follows from the continuity of $\mathbf{T}^\square$ on $W_c^2(I;F)$, and its continuity on $(F_c(I))'$ follows from the continuity of $\mathbf{T}^\square$ on $F_c(I)$. Let us prove this when $q \in L^p_{loc}(I)$, thus $F(I) = L^p(I)$ and $W_c^2(I;F) = W_c^{2,p}(I)$. The other cases will follow by analogy. From (31) and since $K(t,x) \in C(I\times I) \cap W^{1,1}_{loc}(I\times I)$, it is clear that $\mathbf{T}^\square$ is continuous on $L^p_c(I)$. Moreover, if $\phi \in W_c^{2,p}(I)$, then formula (33) holds, and from this we see that $\mathbf{T}\phi \in W_c^{1,p}(I)$. On the other hand, if $\phi, \psi \in W_c^{2,p}(I)$, then integration by parts gives

$$\begin{aligned}\int_I (T^\square\phi)'\psi' dx &= -\int_I (T^\square\phi)\psi'' dx = -\int_I \phi T\psi'' dx = \\ &= -\int_I \phi\left\{(T\psi)'' - qT\psi\right\} dx = -\int_I \left\{T^\square(\phi'' - q\phi)\right\}\psi dx,\end{aligned}$$

where we have used the fact that the functions $\mathbf{T}^\square\phi$ and $\mathbf{T}^\square\psi$ have compact support and also the transmutation property of $\mathbf{T}$ on $W_c^{2,1}(I)$. Thus we have shown that $\mathbf{T}^\square\phi \in W_c^{2,p}(I)$ and

$$(\mathbf{T}^\square\phi)'' = \mathbf{T}^\square(\phi'' - q\phi),\ \phi \in W_c^{2,p}(I). \tag{34}$$

Now, by (33) and (34) we see that $\mathbf{T}^\square\varphi_n \to \mathbf{T}^\square\varphi$ in $W_c^{2,p}(I)$, whenever $\varphi_n \to \varphi$ in $W_c^{2,p}(I)$, as $n\to\infty$. This ensures the continuity of $\mathbf{T}^\square$ on $W_c^{2,p}(I)$.

By analogy with the proof of Theorem 26, we arrive at the invertibility of $\mathbf{T}$.

Lastly, let us prove the transmutation property of $\mathbf{T}$. If $u \in (F_c(I))' \subset (W_c^2(I;F))'$, then u'', $\mathbf{T}u''$, $q\mathbf{T}u$, $(\mathbf{T}u)'' \in (W_c^2(I;F))'$ and using (34) we see that

$$\left\langle (\mathbf{T}u)'' - q\mathbf{T}u, \phi\right\rangle = \left\langle u, \mathbf{T}^\square(\phi'' - q\phi)\right\rangle = \left\langle u, (\mathbf{T}^\square\phi)''\right\rangle = \langle \mathbf{T}u'', \phi\rangle$$

holds for all $\phi \in W_c^{2,p}(I)$. ■

Remark 31 *Let $q \in F_{loc}(I)$. Reasoning as in Corollary 27, we can conclude that the equation $u'' = qu$, $u \in (F_c(I))'$ has only weak solutions $u \in W^{2,1}(I)$. Thus, as before, we will be able to consider standard transmutation operators. Moreover, reasoning as in Proposition 29, we see that an s-transmutation operator on $(F_c(I))'$ that maps continuously $(F_c(I))'$ to $(W_c^1(I;F))'$ must have the form $\mathcal{T} = c_1\mathbf{T}P_+ + c_2\mathbf{T}AP_+ + c_3\mathbf{T}\frac{d}{dx}P_- + c_4\mathbf{T}P_-$ where the c_i are constants. Moreover, $\mathcal{T}$ maps $(F_c(I))'$ to itself continuously if and only if $c_3 = 0$.*

7 Conclusions and generalizations

A method to compute the general formula of the s-transmutation operators corresponding to $L = \frac{d^2}{dx^2} - q(x)$ has been presented. On the one hand, the method is based on fairly simple formulas, which allows one to compute a fundamental system of s-transmutation operators when one single (one-to-one) s-transmutation operator is known in closed form. On the other hand, with the aid of classical results we established that a particular s-transmutation operator can be represented in the form of a Volterra integral operator of the second kind.

Most of the results of Sections 3 and 4 can be extended to the second-order linear differential operator $L_1 = a_2(x)\frac{d^2}{dx^2} + a_1(x)\frac{d}{dx} + q(x)$ with smooth coefficients. This includes the explicit formulas leading to the construction of new s-transmutations and then to the general formula of the s-transmutations corresponding to L_1, once the closed form of one single s-transmutation corresponding to L_1 is known. If $a_2(x) \equiv 1$, then a closed form for such an s-transmutation can be obtained following [26, Chap. 1, Th. 7.1] and if $L_1 = \frac{d}{dx}p\frac{d}{dx} + q(x)$, by following [20].

It would also be interesting to develop this method for higher order linear differential operators. However, apart from the results of [1] and [15], there are, as far as our knowledge extends, no results known about the existence of transmutation operators for higher orders.

References

[1] Begehr H., and Gilbert R., *Transformations, Transmutations and Kernel Functions*, vols. 1–2. Pitman, London, 1992.

[2] Barros-Neto J., *An Introduction to the Theory of Distributions*. Marcel Dekker, New York, 1973.

[3] Bers L., *Theory of Pseudo-Analytic Functions*. New York University, 1952.

[4] Blancarte H., Campos H. M., and Khmelnytskaya K. V., Spectral parameter power series method for discontinuous coefficients. *Mathematical Methods in the Applied Sciences* 38(10), 2000–2011, 2015.

[5] Brezis H., *Functional Analysis, Sobolev Spaces and Partial Differential Equations*. Springer-Verlag, Berlin, 2011.

[6] Campos H. M., *Bicomplex Pseudoanalytic Functions and Applications in Constructive Methods for Solving Boundary Value Problems*. PhD thesis, Cinvestav, Mexico City, 2012.

[7] Campos H. M., Kravchenko V. V., and Mendez L. M., Complete families of solutions for the Dirac equation: An application of bicomplex pseudoanalytic function theory and transmutation operators. *Adv. Appl. Clifford Algebr.* 22(3) 577–594, 2012.

[8] Campos H. M., Kravchenko V. V., and Torba S. M., Transmutations, L-bases and complete families of solutions of the stationary Schrödinger equation in the plane. *Journal of Mathematical Analysis and Applications* 389(2), 1222–1238, 2012.

[9] Carroll R., *Transmutation Theory and applications*. North-Holland, Amsterdam, 1985.

[10] Duistermaat, J. J., and Kolk, J. A. C., *Distributions: Theory and Applications*. Birkhäuser, New York, 2010.

[11] Fage M. K., and Nagnibida N. I., *The Problem of the Equivalence of Ordinary Linear Differential Operators*. Nauka, Novosibirsk, USSR, 1987 (in Russian).

[12] Grubb G., *Distributions and Operators*. Springer-Verlag, Berlin, 2009.

[13] Khmelnytskaya K. V., Kravchenko V. V., and Torba S. M., Modulated electromagnetic fields in inhomogeneous media, hyperbolic pseudoanalytic functions and transmutations, submitted, available at arXiv:1410.4873.

[14] Khmelnytskaya K. V., Kravchenko V. V., Torba S. M., and Tremblay S., Wave polynomials and Cauchy's problem for the Klein-Gordon equation. *J. Math. Anal. Appl.* 399, 191–212, 2013.

[15] Kiryakova V., Transmutation method for solving hyper-Bessel differential equations based on the Poisson–Dimovski transformation. *Fractional Calculus and Applied Analysis* 11(3), 2008.

[16] Kolmogorov A. N., and Fomin S. V., *Introductory Real Analysis*. Dover, New York, 1975.

[17] Kravchenko V. V., A representation for solutions of the Sturm-Liouville equation. *Complex Variables and Elliptic Equations* 53(8), 775–789, 2008.

[18] Kravchenko V. V., *Applied Pseudoanalytic Function Theory*. Birkhäuser, Basel, 2009.

[19] Kravchenko V. V., and Porter R. M., Spectral parameter power series for Sturm–Liouville problems. *Mathematical Methods in the Applied Sciences* 33, 459–468, 2010.

[20] Kravchenko V. V., Morelos S., and Torba S. M., Liouville transformation, analytic approximation of transmutation operators and solution of spectral problems. Submitted, available at arXiv:1412.5237.

[21] Kravchenko V. V., and Torba S. M., Construction of transmutation operators and hyperbolic pseudoanalytic functions. *Complex Analysis and Operator Theory* 9(2), 379–429, 2015.

[22] Kravchenko V. V., and Torba S. M., Transmutations for Darboux transformed operators with applications. *Journal of Physics A: Mathematical and Theoretical* 45(7), 075201, 2012.

[23] Kravchenko V. V., and Torba S. M., Transmutations and spectral parameter power series in eigenvalue problems. In: *Operator Theory, Pseudo-Differential Equations, and Mathematical Physics*, pp. 209–238, Springer-Verlag, Berlin, 2013.

[24] Kravchenko V. V., and Torba S. M., Analytic approximation of transmutation operators and applications to highly accurate solution of spectral problems, *J. Comput. Appl. Math.* 275, 1–26, 2015.

[25] Levitan B. M., *Inverse Sturm–Liouville Problems*. VSP, Zeist, 1987.

[26] Lions J.-L., Opérateurs de Delsarte et problèmes mixtes. *Bulletin de la S. M. de France* 84, 9–95, 1956.

[27] Marchenko V. A., *Sturm–Liouville Operators and Applications*. Birkhäuser, Basel, 1986.

[28] Treves F., *Topological Vector Spaces, Distributions, and Kernels*. Academic Press, New York, 1967.